



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Climate-smart soils

Citation for published version:

Paustian, K, Lehmann, J, Ogle, S, Reay, D, Robertson, GP & Smith, P 2016, 'Climate-smart soils', *Nature*, vol. 532, no. 7597, pp. 49–57. <https://doi.org/10.1038/nature17174>

Digital Object Identifier (DOI):

[10.1038/nature17174](https://doi.org/10.1038/nature17174)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Nature

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



‘Climate-smart’ soils: a new management paradigm for global agriculture

Keith Paustian^{1,2*}, Johannes Lehmann³, Stephen Ogle^{2,4}, David Reay⁵, G. Philip Robertson⁶ and Pete Smith⁷

¹ Dept. of Soil and Crop Sciences, Colorado State University, Fort Collins, CO, USA

² Natural Resource Ecology Lab, Colorado State University, Fort Collins, CO, USA

³ Dept. of Soil and Crop Sciences, Cornell University

⁴ Dept. of Ecosystem Science and Sustainability, Colorado State University, Fort Collins, CO, USA

⁵ School of Geosciences, University of Edinburgh, Edinburgh, Scotland, UK

⁶ W.K. Kellogg Biological Station and Dept of Plant, Soil and Microbial Sciences, Michigan State University, E. Lansing, MI, USA

⁷ Institute of Biological and Environmental Sciences, University of Aberdeen, Aberdeen, Scotland, UK

* Corresponding author

Preface

Soils are integral to the function of all terrestrial ecosystems and for sustaining food and fibre production. An overlooked aspect of soils is their potential to mitigate greenhouse gas (GHG) emissions. Although proven practices exist, implementation of soil-based GHG mitigation activities are early-stage and accurately quantifying emissions and reductions remains a significant challenge. Emerging research and information technology developments provide the

potential for broader inclusion of soils in GHG policies. We highlight ‘state-of-the-art’ soil GHG research, summarize mitigation practices and potentials, identify gaps in data and understanding and suggest ways to close gaps through new research, technology and collaboration.

Introduction

Evidence points to agriculture as the first instance of human-caused increases in greenhouse gases (GHGs), several thousand years ago¹. Agriculture and associated land use change remain a source for all three major biogenic GHGs -- carbon dioxide (CO₂), methane (CH₄), and nitrous oxide (N₂O). Land use contributes ~25% of total global anthropogenic GHG emissions: 10-14% directly from agricultural production, mainly via GHG emissions from soils and livestock management, and another 12-17% from land cover change, including deforestation^{2,3}. While soils contribute a major share (37%; mainly as N₂O and CH₄) of agricultural emissions³, improved soil management can substantially reduce these emissions and sequester some of the CO₂ removed from the atmosphere by plants, as carbon (C) in soil organic matter (in this paper, our discussion of soil C refers solely to organic C). In addition to decreasing GHG emissions and sequestering C, wise soil management that increases organic matter and tightens the soil nitrogen (N) cycle can yield powerful synergies, such as enhanced fertility and productivity, increased soil biodiversity, reduced erosion, runoff and water pollution, and can help buffer crop and pasture systems against the impacts of climate change⁴.

The inclusion of soil-centric mitigation projects within GHG offset markets⁵ and new initiatives to market ‘low-carbon’ products⁶ indicate a growing role for agricultural GHG mitigation⁷.

Moreover, interest in developing aggressive soil C sequestration strategies has been heightened by recent IPCC assessments, which project that substantial terrestrial C sinks will be needed to supplement large cuts in GHG emissions to achieve GHG stabilization levels of 450ppm CO₂ equivalent or below, consistent with the goal of <2° C mean global temperature increase⁸. Soil C sequestration is one of a few strategies that could be applied at large scales⁸ and potentially at low cost; as an example, the French government is proposing a plan to increase soil C concentration in a large portion of agricultural soils globally, by 0.4% per year, producing a C sink increase of 1.2 Pg C yr⁻¹[9].

An extensive body of field, laboratory and modelling research over many decades demonstrates that improved land use and management practices can reduce soil GHG emissions and increase soil C stocks. However, implementing effective soil-based GHG mitigation strategies at scale will require capacity to measure and monitor GHG reductions with acceptable accuracy, quantifiable uncertainty and at relatively low cost. Targeted research to improve predictive models, expanded observational networks to support model validation and uncertainty bounds, ‘Big Data’ approaches to integrate land use, management and environmental drivers, and technologies to actively engage with land users at the grass-roots, are key elements to realizing the potential GHG mitigation from ‘climate smart’ agricultural soils.

Process controls and mitigation practices

Soil C sequestration via improved management

Soils constitute the largest terrestrial organic C pool (ca. 1500 Pg C to 1 m depth; 2400 Pg C to 2 m depth¹⁰), which is three times the amount of CO₂ currently in the atmosphere (~830 Pg C) and

240 times current annual fossil fuel emissions ($\sim 10 \text{ Pg}$)⁸. Thus, increasing net soil C storage by even a few percent represents a significant C sink potential.

Proximal controls on the soil C balance include the rate of C addition as plant residue, manure or other organic waste, less the rate of C loss (*via* decomposition); hence, C stocks can be increased by increasing organic matter inputs or by reducing decomposition rates (e.g., by reducing soil disturbance), or both, leading to net removal of C from the atmosphere¹¹. However, soil C accrual rates decrease over time as stocks approach a new equilibrium. Thus net CO₂ removals are of limited duration, often attenuating after 2-3 decades¹².

Unmanaged forests and grasslands typically allocate a large fraction of their biomass production belowground and their soils are relatively undisturbed; accordingly, native ecosystems usually support significantly higher soil C stocks than their agricultural counterparts, and soil C loss (typically 0.5 to $>2 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$) following land conversion to cropland has been extensively documented^{13,14}. Total losses once the soil approaches a new equilibrium are typically $\sim 30\text{-}50\%$ of topsoil (e.g. $0\text{-}30 \text{ cm}$) C stocks¹⁴. Hence, avoided conversion and degradation of native ecosystems is a strong mitigation alternative. Conversely, restoration of marginal or degraded lands to perennial forest or grassland increases soil C storage (Fig. 1), although usually at a slower rate than the original conversion losses^{15,16}. Restoring wetlands that have been drained for agricultural use reduces ongoing decomposition losses, which can be as high as $5\text{-}20 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$ [17], and can also restore C sequestration (Fig. 1), though methane emissions may increase^{18,19}. Land use conversions may, however, conflict with agricultural production and food security objectives, entailing the need for a broad-based accounting of net GHG implications²⁰.

[Fig 1 about here]

In general, soil C sequestration rates on land maintained in agricultural use are less than for land restoration/conversion, and vary on the order of 0.1 to 1 Mg C ha⁻¹ yr⁻¹, as a function of land use history, soil/climate conditions, and the combination of management practices applied^{2,14}.

Practices that increase C inputs include (i) improved varieties or species with greater root mass to deposit C in deeper layers where turnover is slower²¹, (ii) adopting crop rotations that provide greater C inputs²², (iii) more residue retention²³, and (iv) cover crops during fallow periods to provide year-round C inputs (Fig. 1).^{22,24} Cover crops can also reduce nutrient losses, including nitrate that is otherwise converted to N₂O in riparian areas and waterways²⁵ – an example of synergy between practices that sequester C and also tighten the N cycle to limit emissions of N₂O. Other practices to increase C inputs include irrigation in water-limited systems¹⁸ and additional fertilizer input to increase productivity in low-yielding, nutrient deficient systems (Fig. 1)²⁶. Although additional nutrient and water inputs to boost yields may increase non-CO₂ emissions²⁷, the emissions intensity of the system (GHG emissions per unit yield) may decline, providing a global benefit if the yield increase avoids land conversion for agriculture elsewhere^{20,22}.

Some croplands can sequester C through less intensive tillage, particularly zero tillage¹⁴, due to less disruption of soil aggregate structure²⁸. Some authors have argued that benefits are small because increased C content in surface horizons are offset by C losses deeper in the profile²⁹,

although others have noted that the larger variability in sub-surface horizons and lack of statistical power in existing studies makes such conclusions questionable³⁰.

A change from annual to perennial crops typically increases belowground C inputs (and soil disturbance is reduced), leading to C sequestration¹⁵. In grasslands, soil C sequestration can be increased through optimal stocking/grazing density³¹. Improved management in fire-prone ecosystems *via* fire prevention or prescribed burning can also increase C sequestration³².

Key knowledge gaps that affect our understanding of soil C sequestration processes and management options to implement them include questions about the differential temperature sensitivity of C turnover among SOM fractions³³, interactions among organic matter chemistry, mineral surface interactions and C saturation³⁴⁻³⁶, and subsoil (> 30 cm) SOM accretion, turnover and stabilization³⁷. Landscape processes, particularly the impact of erosion and lateral transport of C in sediments, contribute additional uncertainty on net sequestration occurring at a specific location³⁸. And emerging evidence that stabilized SOM is of microbial rather than direct plant origin^{34,39} may offer a potential to manipulate the soil-plant microbiome to enhance C sequestration in the rhizosphere.

Soil C sequestration via exogenous C inputs

Addition of plant-derived C from external (i.e., offsite) sources such as composts or biochar can increase soil C stocks, and may result in net CO₂ removals from the atmosphere (Fig. 1). Both compost and biochar are more slowly decomposed compared to fresh plant residues, with composts typically having mean residence times several-fold greater than un-composted organic

matter⁴⁰, and biochar mineralizes 10-100 times slower than uncharred biomass⁴¹. Thus a large fraction of added C — particularly for biochar — can be retained in the soil over several decades or longer, although residence times vary depending on the amendment type, nutrient content and soil conditions³⁵ (e.g. moisture, temperature, texture).

However, because the organic matter originates from outside the ecosystem ‘boundary’, a broader life-cycle assessment approach is needed, that considers GHG impacts of: (i) offsite biomass removal, transport, and processing, (ii) alternative end uses of the biomass, (iii) interactions with other soil GHG-producing processes, and (iv) synergies between these soil amendments and the fixation and retention of *in situ* plant-derived C^{42,43}. In many cases, net life-cycle emissions will largely depend on whether the biomass used as a soil amendment would have otherwise been burnt (either for fuel, thereby offsetting fossil fuel use, or as waste disposal), added to a landfill, or left in place as living biomass or detritus^{42,43}.

While slower mineralization of the amendment is an important determinant of net mitigation impact, effects on other soil emissions cannot be neglected. Mineralization of existing soil C in response to amendments (often referred to as ‘priming’⁴⁴) has often been observed immediately following biochar addition, but priming usually declines, sometimes becoming negative (i.e., inhibiting *in situ* soil C decomposition), over time^{45,46}. Analogous time dependence of soil N₂O and CH₄ emissions has not received sufficient attention⁴⁰. Increased plant growth in amended soils and the resultant feedbacks to soil C can make up a large proportion of the soil-based GHG balance^{40,47} and these feedbacks may be especially important for more persistent amendments, because of the longer duration of any effects.

Soil management to reduce N₂O emissions

Arable soils emit more N₂O to the atmosphere than any other anthropogenic source^{2,18}; some 4.2 Tg of a global anthropogenic flux of 8.1 Tg N₂O-N yr⁻¹. Reducing this flux represents a significant mitigation opportunity, particularly since N₂O is often the major source of radiative forcing in intensively managed cropland. Better N management to reduce emissions would also ameliorate other environmental problems such as nitrate pollution of ground and surface waters caused by excess reactive N in agroecosystems (Fig. 1).

N₂O is produced in soils by microbial activity – mainly nitrification and denitrification – which occur readily when stimulated by the abundant N that cycles rapidly in virtually all agroecosystems. During nitrification, ammonium added as fertilizer, fixed from the atmosphere by legumes, or mineralized from soil organic matter, crop residue, or other inputs is oxidized to nitrite and eventually to nitrate in a series of reactions that can also produce N₂O. Likewise, when denitrifiers use nitrate as an electron acceptor when soil oxygen is low, N₂O is an intermediate product that can readily escape to the atmosphere.

Arable soils managed to support high crop productivity have the capacity to produce large quantities of N₂O, and fluxes are directly related to N inputs. On average, about 1% of the N applied to cropland is directly emitted as N₂O⁴⁸, which is the basis for estimating emissions using default IPCC methods¹⁷. However, recent evidence suggests that this value is too high for crops that are under-fertilized and too low for crops that are fertilized liberally²⁷. When crops compete with microbes for available N, N₂O fluxes are lower. In addition to direct in-field

emissions, high N applications cause N losses from leaching and volatilization that contribute to ‘indirect’ N₂O emissions, downstream/downwind from the field⁴⁹.

Since N₂O has no significant terrestrial sink, abatement is best achieved by attenuating known sources of N₂O emissions, by altering the environmental factors that affect N₂O production (soil N, oxygen, and C) or by biochemically inhibiting conversion pathways using soil additives. For example, nitrification can be inhibited with commercial additives such as nitrapyrin and dicyandiamide, which slow ammonium oxidation, and field experiments suggest that inhibitors can reduce N₂O fluxes up to 40% in some soils, although other soils show little reduction and more research is needed to understand variable site-level responses⁵⁰. Likewise, tillage and water management can affect N₂O fluxes by altering the soil microenvironment^{51,52}.

Another means for reducing N₂O emissions from arable soils is more precise N management to minimize excess N not used by the crop, while maintaining sustainable high yields. Fertilized crops typically take up less than 50% of the N applied; the remainder is available for loss. By one recent study⁵³, corn farmers in the U.S. Midwest could reduce N₂O loss by 50% with more conservative fertilizer practices. Nitrogen conservation can be achieved by: (1) better matching application rates of N to crop needs using advanced statistical and quantitative modelling; (2) applying fertilizer at variable rates across a field based on natural patterns of soil fertility, or within the root zone rather than broadcast on the soil surface; and (3) applying fertilizer close to when the crop can use it, such as several weeks after planting, or adding it earlier but using slow-release coatings to delay its dissolution⁴⁹.

High temporal and spatial variability make predictions of changes in N₂O fluxes in response to management surprisingly difficult. Particularly lacking are empirical data for multi-intervention strategies that may interact in unexpected ways. Aligned to this paucity are gaps in our understanding of how N cycling and net N₂O flux in managed soils will respond to future climate change⁵⁴. The limited number of field manipulation studies to date indicate that changing temperature and precipitation patterns may have large and strongly-coupled effects on net N₂O emissions⁵⁵, yet our understanding of the processes that underpin these effects and their robust representation in models is far from complete.

Soil management to reduce CH₄ emissions

More than one-third (>200 Tg yr⁻¹)⁸ of global methane (CH₄) emissions occur through the microbial breakdown of organic compounds in soils under anaerobic conditions⁵⁶. As such, wetlands (177-284 Tg yr⁻¹) and rice cultivation (33-40 Tg yr⁻¹)⁸ represent the largest soil-mediated sources of CH₄ globally. In contrast, well-aerated soils act as sinks for CH₄ (estimated at ~ 30 Tg yr⁻¹) from the atmosphere *via* CH₄ oxidation, the bulk of this net sink being in unmanaged upland and forest soils⁵⁷.

Key determinants of soil CH₄ fluxes include aeration, substrate availability, temperature and N inputs⁵⁸; therefore, soil management can radically alter CH₄ fluxes. For example, in most soils, conversion to agriculture severely restricts CH₄ oxidation, related to the suppression of methanotrophs by accelerated N cycling⁵⁹. In flooded rice, alterations in drainage regimes and organic residue incorporation could reduce emissions by ~ 25% or 7.6 Tg CH₄ yr⁻¹ globally¹⁸,

although cycles of wetting and drying of soils may also enhance N₂O production⁶⁰ and soil C mineralisation⁶¹, thereby reducing the net mitigation effect.

With global rice production projected to expand by ~40% between 2000-2023 [62], the potential for further GHG mitigation via soil management appears large, although the global distribution and diverse nature of rice production systems – including irrigated, rain-fed and deepwater – present challenges to developing effective mitigation strategies. For longer-term (>20 year) projections, climate change and land-atmosphere interactions become increasingly important, with changes in N inputs, temperature, precipitation and atmospheric CO₂ concentration all likely to affect net CH₄ fluxes from soils⁶³.

This uncertainty highlights important gaps in understanding key processes and their underlying controls. The restoration of soil CH₄ uptake following agricultural conversion, for example, appears related to methanotroph community diversity⁶⁴, about which we know too little. Likewise the abatement of CH₄ generation in rice rhizospheres is related to C compounds exuded by roots, such that CH₄ mitigation might be achieved through further rice breeding and genetics⁶⁵. Limited availability of field-scale CH₄ flux data means a greater reliance on regionally-averaged emission factors and extrapolation from mesocosm and laboratory incubations¹⁷, and thus less site and condition specificity in modelling fluxes. Importantly, establishing the net climate forcing effects of any intervention is a prime target for future soil management research.

[Fig 2. about here]

Global potential for soil GHG mitigation

How significant, in total, is this large, varied set of land use and management practices as a GHG mitigation strategy? One of the challenges in answering this question is to distinguish between what is technically feasible and what might be achieved given economic, social and policy constraints. A comprehensive global analysis of agricultural-related practices by Smith et al.¹⁸ combined climate-stratified modelling of emission reductions and soil C sequestration with economic and land use change models to estimate mitigation potential as a function of varying 'C prices' (reflecting social incentive to pay for mitigation). They estimated total soil GHG mitigation potential ranging from 5.3 Pg CO₂eq yr⁻¹ (absent economic constraints) to 1.5 Pg CO₂eq yr⁻¹ at the lowest specified C price (\$20 per Mg CO₂eq). Average rates for the majority of management interventions are modest, < 1 Mg CO₂eq ha⁻¹ yr⁻¹. Thus, achieving globally significant GHG reductions requires a substantial proportion of the agricultural land-base (Fig. 2). Although the economic and management constraints on biochar additions (not assessed by Smith et al.¹⁸) are less well known, Woelf et al.⁶⁶ estimated a global technical potential of 1-1.8 Pg CO₂eq yr⁻¹ (Fig. 2).

A more unconventional intervention that has been proposed is the development of crops with larger, deeper root systems, hence increasing plant C inputs and soil C sinks^{21,67}. Increasing root biomass and selecting for root architectures that store more C in soils has not previously been an objective for crop breeders, although most crops have sufficient genetic plasticity to substantially alter root characteristics⁶⁸ and selection aimed at improved root adaptation to soil acidity, hypoxia and nutrient limitations could yield greater root C inputs as well as increased crop yields

⁶⁷. Greater root C inputs is well-recognized as a main reason for the higher soil C stocks maintained under perennial grasses compared to annual crops ¹⁵. Although there are no published estimates of the global C sink potential for ‘root enhancement’ of annual crop species, as a first-order estimate, a sustained increase in root C inputs might add ~1 Pg CO₂eq yr⁻¹ or more if applied over a large portion on global cropland area (Fig 2).

Hence, the overall mitigation potential of existing (and potential future) soil management practices could be as high as ~8 Pg CO₂eq yr⁻¹. How much is achievable will depend heavily on the effectiveness of implementation strategies and socioeconomic and policy constraints. A key strength is that a variety of practices can often be implemented on the same land area, to leverage synergies, while avoiding offsetting effects for different gases (Fig. 1). But regardless of which combination of management interventions are pursued, effective policies, that incentivize land managers to adopt them, will be needed. A common thread across implementation strategies is the role for strong science-based metrics to measure and monitor performance.

Implementation of mitigation practices

Relative to many other GHG source categories, agricultural soil GHG mitigation presents particular challenges. Rates on an individual land parcel are often low, but vast areas of land are devoted to agriculture globally, and the implementers of mitigation practices – the people using the land – number in the billions. Thus engaging a significant number of these people is a massive undertaking in itself. Furthermore, agricultural soil GHG emissions are challenging to quantify due to their dispersed and variable nature and the multiplicity of controlling factors – operating across heterogeneous landscapes. Direct measurement of fluxes requires specialized

personnel and equipment, normally limited to research environments, and hence not feasible for most mitigation projects. Model-based methods, in which emission rates are quantified as a function of location, environmental conditions and management, provide a more feasible approach^{52,69,70}. Process-based models, which dynamically simulate mechanisms and controls on fluxes as a function of climatic and soil variables and management practices, and empirical models based on statistical analysis of field-measured flux rates, represent differing but complementary approaches. In general, model-based quantification systems enable monitoring to focus on practice performance and thus dramatically reduce transaction costs for implementing mitigation policies⁶⁹.

[Box 1 about here]

Several implementation strategies for soil GHG mitigation exist (see Box 1), all of which require robust quantification and monitoring technologies. Those requiring the most rigorous methods involve offset projects participating in cap-and-trade markets, in which land managers are directly compensated for achieving emission reductions. Other market-linked strategies, such as ‘green labeling’ systems for agricultural products, will also require rigorous yet easy to use GHG quantification tools, enabling agricultural producers to meet standards set by product distributors and accepted by consumers^{6,71}.

Within the voluntary C offset market space, there are a growing number of projects that include soil GHG mitigation components⁵. Several large projects focus on preventing land conversion (i.e., from forest and grassland), thus avoiding large CO₂ emissions from soils and liquidated

biomass C stocks. Relatively simple empirical models supplemented with field measurements are commonly used for avoided land conversion projects. For more complex land use projects, empirical models are less suited to capture interactions across multiple emission sources, and may over- or under-credit projects where a practice has an influence on multiple emission sources. There are relatively fewer projects targeting GHG mitigation on existing agricultural lands, involving a broader suite of soil management practices, and early pilot-phase N₂O and CH₄ reduction projects are only now being developed^{5,52}. Here, accurately quantifying C sequestration and/or emission reductions is more challenging due to lower rates of change relative to baseline conditions, thus requiring more sophisticated models and supporting research infrastructure (Fig. 3).

Another challenge for projects on existing agricultural lands is obtaining and processing the management activity data. For example, the Kenya Agriculture Carbon Project (KACP) involves a total of 60,000 individual small-holder farmers⁷². In contrast to projects involving major land cover changes, where remote sensing can provide much of the activity monitoring (e.g., retention of forested land over time), such options are poorly-suited for monitoring crop type, fertilizer, residue and water management, and organic matter amendments⁷³; for such practices the best source of information are the land managers themselves (Fig. 3).

Thus another option is to engage land managers as information providers. Examples of this approach are the Cool Farm Tool⁷¹, being used by farmers participating in low C supply chain management, and the COMET-Farm tool, which allows farmers to compute full farm-scale GHG budgets, for support of government-sponsored conservation initiatives and participation in

mitigation projects⁷⁴. Both tools provide web-based interfaces designed for non-specialists to enter land management information; Cool Farm utilizes empirical emission factor-type models, while COMET-Farm incorporates both empirical and process-based models. Such systems can be used to integrate local knowledge on management practices with detailed soil and climate maps, remote sensing and sophisticated models for emission calculations. Soon much of this functionality could be deployed in mobile applications (Fig. 3), which would be particularly advantageous in developing countries where existing infrastructure to collect and manage land use data is weak⁷⁵.

[Fig. 3 about here]

Quantifying uncertainties

Inventories of soil C stock changes and net GHG fluxes using process-based models will always have uncertainty due to lack of process understanding, inadequate parameterization, and limitations associated with model inputs⁷⁶ (e.g., weather, management and soils data). Empirical models generally rely on statistical analyses of measurement data to produce emission factors, along with an estimated uncertainty¹⁴. However, empirical models can be biased if measurements do not fully reflect the conditions for the agroecosystems in the project. Even with the limitations in process-based understanding, process-based models are likely to provide the most robust framework for estimating soil C stock and GHG flux changes in climate smart agriculture programs⁷⁷.

Monitoring, reporting and verification (MRV) systems are a key element in a climate smart agricultural program. While MRV systems place different levels of importance on uncertainty depending on program type (see Box 1)⁷⁸, discounting payments based on the level of uncertainty is likely to be part of programs with financial incentives, such as cap-and-trade. Discounting encourages monitoring efforts to reduce uncertainty over time¹⁷. If discounting payments for C sequestration and emission reduction practices with larger uncertainty is adopted in climate smart agriculture programs, then more advanced methods with process-based models will likely emerge as the preferred method due to less uncertainty. For example, uncertainty was reduced by 24% when predicting national-scale C stock changes in the United States with process-based models compared to empirically-derived factors⁷⁶.

Another consideration is that uncertainties in estimating C stock and GHG emissions with process-based models are considerably larger for reporting by single individuals, particularly if the amount of change on an individual farm is small⁷⁶. Aggregation of many farms into larger projects will reduce uncertainties, which could be a viable approach for managing uncertainty and reducing discounting of incentive payments.

Verification is an independent evaluation of estimated emissions intended to provide confidence that the reported results are correct, but in practice, the requirements for verification are highly variable across different GHG mitigation efforts, from essentially no requirements to annual evaluations⁷⁸. Verification typically focuses on the accuracy of the estimates, and possibly the most stringent approach is an independent set of measurements. Although independent data may be less favored in terms of costs relative to alternatives, such as expert judgement⁷⁸, soil

monitoring networks deployed at national or regional scales could produce independent data for evaluating model-based assessments of soil C stock changes and GHG emissions⁷⁹ and for model bias adjustment, using empirically-based methods⁸⁰.

Another approach to verification is to use atmospheric observations of trace gas concentrations and inverse modeling to estimate fluxes between the atmosphere and land surface^{81,82}. This ‘top-down’ modeling, utilizing a network of tower-based observations of CO₂ concentrations, was used to verify ‘bottom-up’ inventory modeling based on observed management activities, in the largely agricultural region of the central United States^{83,84}. Since atmospheric observations integrate all CO₂ fluxes in the region, the inventory included a full assessment of all sources and sinks. However, even with the fully integrated CO₂ flux, it is possible to statistically disaggregate individual sources as part of the analysis, such as contributions from soil C pools to the regional flux⁸⁵. Satellite-based measurements are providing a new source of atmospheric trace gas data that can be used to estimate land surface fluxes with inverse modeling frameworks^{86,87}. While atmospheric observations and satellite imagery may become a standard for verifying regional inventories in the future, the methods need further testing in the near term before deploying operational systems.

Conclusions and way forward

Climate change and GHG mitigation require an ‘all of the above’ approach⁸⁸, where all reduction measures that are feasible, cost-effective and environmentally sustainable should be pursued. For soils, a variety of management practices and technologies are known to reduce emissions and promote C sequestration, most of which also provide environmental co-benefits. Impediments to

more aggressively implementing agricultural soil GHG mitigation strategies to date are primarily the feasibility of cost-effectively quantifying and verifying soil mitigation activities⁸⁹.

Overcoming these barriers therefore translates into: i) increasing the acceptance of soil management within compliance and voluntary C markets, ii) reducing costs to governments for providing environmental-based subsidies, and iii) meeting demands of consumers for ‘low carbon’ products.

Reducing and managing uncertainties are key to both improved predictive models and decision-support tools and the design of effective policies that promote soil-based GHG mitigation. To advance these efforts, several research and development priorities are apparent (Fig 3). First, support for research site networks of soil flux (N_2O , CH_4) and soil C measurements⁹⁰ encompassing a wide variation in management, as well as ‘on-farm’ soil C monitoring networks⁷⁹ needs to be strengthened, in coordination with basic research (e.g., on SOM stabilization processes, N_2O and CH_4 microbiology, plant-microbe interactions, plant breeding and root phenotyping) to advance process understanding, develop new mitigation practices and fill gaps for underrepresented soil/climate/management systems. High quality data generated from consistent measurement protocols is critical for evaluating and improving models. These efforts may benefit from development of new sensor technologies enabling cheaper and quicker soil measurements⁹¹. While multiple competing models are needed, both to spur innovation and because no single model will be best in all situations, model development will benefit from greater collaboration and cross-model testing among developers, moving towards a more open-source, community development approach⁹². Large geospatial databases of soil biophysical properties and climate variables are critical to accurately quantify soil processes across the

landscape (Fig. 3). High resolution soil maps exist in most developed countries (and increasingly in developing countries⁹³), and if made publically available⁹⁴, would greatly improve capabilities for modeling GHG emission at scale.

Finally, realising the potential for climate change mitigation through global soil management requires understanding cultural, political and socioeconomic contexts, and the ways in which widespread, sustained changes in practice can be successfully achieved within it^{95,96}. As such, there needs to be greater level of engagement with the land users themselves, who will be the ones implementing practices that abate GHG emissions and sequester C. Engagement means both education and outreach, highlighting the links between agriculture and GHGs and utilizing innovative strategies⁷⁵ (Fig. 3) to involve stakeholders in gathering and using their local knowledge of how the land is being used now and how it might best be used in the future, establishing a new paradigm for climate-smart soil management.

References

- [1] Ruddiman, W.F. The anthropogenic greenhouse era began thousands of years ago. *Clim. Chang.* **61**, 261-293 (2003).
- [2] Smith P. *et al.* Agriculture, Forestry and Other Land Use (AFOLU), in *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Edenhofer, O. *et al.*). Cambridge University Press, Cambridge, UK and New York, NY, USA (2014).
- [3] Tubiello, F. N., *et al.* The contribution of agriculture, forestry and other land use activities to global warming, 1990–2012. *Glob. Change Biol.* **21**, 2655-2660 (2015).
- [4] Smith, P. Soils and climate change. *Curr. Opin. Environ. Sustainability* **4**, 539–544 (2012).
- [5] Offset credits marketed through Verified Carbon Standard (VCS) <http://www.v-c-s.org/> and American Carbon Registry (ACR), <http://americancarbonregistry.org/>, with project documentation available on the respective websites (Accessed May, 2015).
- [6] Lavallée, S. & Plouffe, S. The ecolabel and sustainable development. *Int. J. Life Cycle Ass.* **9**(6), 349-354 (2004).
- [7] Kahiluoto, H., Smith, P., Moran, D & Olesen, J.E. Enabling food security by verifying agricultural carbon sequestration. *Nature Clim. Change* **4**, 309-311 (2014).
- [8] Ciais, P. *et al.* Carbon and Other Biogeochemical Cycles, in *Climate Change 2013: The Physical Science Basis* (eds Stocker, T. F. *et al.*) (Cambridge Univ. Press, 2013).

- 468 [9] French government initiative to promote soil C sequestration and mitigation of soil GHG
469 emissions in conjunction with the Conference of the Parties (COP) to the UN Framework
470 Convention on Climate Change (UNFCCC) negotiations in December, 2015, by French
471 (<http://agriculture.gouv.fr/join-40-initiative-soils-food-security-and-climate-0>) (Accessed
472 Sept 2015)
- 473 [10] Batjes, N. H. Total carbon and nitrogen in the soils of the world. *Eur. J. Soil Sci.* **47**, 151-163
474 (1996).
- 475 [11] Paustian, K. *et al.* Agricultural soil as a C sink to offset CO₂ emissions. *Soil Use Manag.* **13**,
476 230-244 (1997).
- 477 [12] West, T.O. & Six, J. Considering the influence of sequestration duration and carbon
478 saturation on estimates of soil carbon capacity. *Clim. Change* **80**, 25-41 (2007).
- 479 [13] Davidson, E. A. & Ackerman, I. L. Changes in soil carbon inventories following cultivation
480 of previously untilled soils. *Biogeochemistry* **20**, 161-193 (1993).
- 481 [14] Ogle, S. M., Breidt, F. J. & Paustian, K. Agricultural management impacts on soil organic
482 carbon storage under moist and dry climatic conditions of temperate and tropical regions.
483 *Biogeochemistry* **72**, 87–121 (2005).
- 484 [15] Conant, R. T., Paustian, K. & Elliott, E. T. Grassland management and conversion into
485 grassland: effects on soil carbon. *Ecol. Appl.* **11**, 343-355 (2001)
- 486 [16] Guo, L.B. & Gifford, R.M. Soil carbon stocks and land use change: a meta analysis. *Glob.*
487 *Change Biol.* **8**, 345-360 (2002).

- [17] IPCC. *Intergovernmental Panel on Climate Change Guidelines for National Greenhouse Gas Inventories* (eds Eggleston, S., Buendia, L., Miwa, K., Ngara, T. & Tanabe, K.) (Institute for Global Environmental Strategies, Hayama, Japan, 2006).
- [18] Smith, P., *et al.* Greenhouse gas mitigation in agriculture. *Phil. Trans.R. Soc. B: Biol. Sci.* **363**, 789-813 (2008). **A comprehensive analysis of agricultural GHG emissions and mitigation potentials including estimated C price impacts on mitigation activities.**
- [19] Knox, S. H., *et al.* Agricultural peatland restoration: effects of land-use change on greenhouse gas (CO₂ and CH₄) fluxes in the Sacramento-San Joaquin Delta. *Glob. Change Biol.* **21**, 750-765 (2015).
- [20] Foley, J. *et al.* Solutions for a cultivated planet. *Nature* **478**, 337-342 (2011).
- [21] Kell, D. Large-scale sequestration of atmospheric carbon via plant roots in natural and agricultural ecosystems: why and how. *Phil. Trans. R. Soc. B* **367**, 1589-1597 (2012).
- [22] Burney J. A., *et al.* Greenhouse gas mitigation by agricultural intensification. *P. Natl. Acad. Sci. USA* **107**, 12052–12057 (2010).
- [23] Wilhelm, W.W., Johnson, J.M.F., Hatfield, J.L., Voorhees, W.B., Linden, D.R. Crop and soil productivity response to corn residue removal: A literature review. *Agron. J.* **96**, 1-17 (2004).
- [24] Poeplau, C. & Don, A. Carbon sequestration in agricultural soils via cultivation of cover crops – A meta-analysis. *Agr. Ecosyst. Environ.* **200**, 33-41 (2015). **This paper combines an analysis of globally-distributed field data with simulation modeling to quantify potential soil C increases with adoption of cover crops on previously fallowed soils.**

- 509 [25] Tonitto, C., David, M. B. & Drinkwater, L. E. Replacing bare fallows with cover crops in
510 fertilizer-intensive cropping systems: A meta-analysis of crop yield and N dynamics. *Agr.*
511 *Ecosyst. Environ.* **112**(1), 58-72 (2006).
- 512 [26] Lemke R. L., VandenBygaart, A. J., Campbell, C. A., Lafond, G. P. & Grant, B. Crop
513 residue removal and fertilizer N: Effects on soil organic carbon in a long-term crop rotation
514 experiment on a Udic Boroll. *Agr. Ecosyst. Environ.* **135**, 42–51 (2010).
- 515 [27] Shcherbak, I., Millar, N. & Robertson, G. P. Global meta-analysis of the nonlinear response
516 of soil nitrous oxide (N₂O) emissions to fertilizer nitrogen. *P. Natl. Acad. Sci. USA* **111**,
517 9199-9204 (2014). **This paper shows that N₂O emissions are greater than previously**
518 **thought for soils receiving high rates of N fertilizer.**
- 519 [28] Six, J., Elliot, E. T. & Paustian, K. Soil macroaggregate turnover and microaggregate
520 formation: A mechanism for C sequestration under no-tillage agriculture. *Soil Biol. Biochem.*
521 **32**, 2099-2103 (2000).
- 522 [29] Powlson, D. S., *et al.* Limited potential of no-till agriculture for climate change mitigation,
523 *Nature Clim. Change* **4**, 678-683 (2014).
- 524 [30] Kravchenko, A. N. & Robertson, G. P. Whole-profile soil carbon stocks: The danger of
525 assuming too much from analyses of too little. *Soil Sci. Soc. Am. J.* **75**(1), 235-240 (2010).
- 526 [31] McSherry, M. E. and Ritchie, M. E. Effects of grazing on grassland soil carbon: a global
527 review. *Glob. Change Biol.* **19**, 1347–1357 (2013).
- 528 [32] Scholes, M.J. & Scholes, R.J. Dust unto dust. *Science* **342**, 565-566 (2013).

- 529 [33] Conant, R. T., *et al.* Temperature and soil organic matter decomposition rates - synthesis of
530 current knowledge and a way forward. *Glob. Change Biol.* **17**(11), 3392-3404 (2011).
- 531 [34] Cotrufo, M. F., Wallenstein, M., Boot, C. M., Deneff, K. & Paul, E. A. The Microbial
532 Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition
533 with soil organic matter stabilization: Do labile plant inputs form stable soil organic matter?
534 *Glob. Change Biol.* **19**, 988-995 (2013).
- 535 [35] Schmidt, M. W. I., *et al.* Persistence of soil organic matter as an ecosystem property. *Nature*
536 **478**, 49-56 (2011).
- 537 [36] Stewart, C. E., Plant, A. F., Paustian, K., Conant, R. & Six, J. Soil carbon saturation:
538 Linking concept and measurable carbon pools. *Soil Sci. Soc. Am. J.* **72**, 379-392 (2008).
- 539 [37] Rumpel, C. & Koegel-Knabner, I. Deep soil organic matter – a key but poorly understood
540 component of terrestrial C cycle. *Plant and Soil* **338**, 143-158 (2011).
- 541 [38] Nadeu, E., Gobin, A., Fiener, P., van Waseael, B. & van Oost, K. Modelling the impact of
542 agricultural management on soil carbon stocks at the regional scale: the role of lateral fluxes.
543 *Glob. Change Biol.* **21**, 3181-3192 (2015).
- 544 [39] Grandy, A. S. & Neff, J. C. Molecular C dynamics downstream: the biochemical
545 decomposition sequence and its impact on soil organic matter structure and function. *Science*
546 *of the Total Environment* **404**, 297-307 (2008).
- 547 [40] Ryals, R., Hartman, M. D., Parton, W. J., DeLonge, M. & Silver W. L. Long-term climate
548 change mitigation potential with organic matter management on grasslands. *Ecol. Appl.* **25**,
549 531–545 (2015). **Field experiments and modeling were used to show reductions in net**

GHG emissions from grazed annual grasslands using composted green waste applications.

- [41] Lehmann J. *et al.* Persistence of biochar in soil, in *Biochar for Environmental Management: Science, Technology and Implementation* (eds Lehmann, J. & Joseph, S.) 235-282 (Taylor and Francis, London, 2015).
- [42] Roberts, K., Gloy, B., Joseph, S., Scott, N. & Lehmann, J. Life cycle assessment of biochar systems: Estimating the energetic, economic and climate change potential. *Envir. Sci. Technol.* **44**, 827–833 (2010).
- [43] DeLonge, M. S., Ryals, R. & Silver, W. L. A lifecycle model to evaluate carbon sequestration potential and greenhouse gas dynamics of managed grasslands. *Ecosystems* **16**, 962-979 (2013).
- [44] Kuzyakov, Y. Priming effects: Interactions between living and dead organic matter. *Soil Biol. Biochem.* **42**, 1363–1371 (2010).
- [45] Zimmerman, A., Gao, B. & Ahn, M. Y. Positive and negative mineralization priming effects among a variety of biochar-amended soils. *Soil Biol. Biochem.* **43**, 1169-1179 (2011).
- [46] Whitman, T., Zhu, Z. & Lehmann, J. Carbon mineralizability determines interactive effects on mineralization of pyrogenic organic matter and soil organic carbon. *Environ. Sci. Tech* **48**, 13727–13734 (2014).
- [47] Whitman, T., Nicholson, C. F., Torres, D., & Lehmann, J. Climate change impact of biochar cook stoves in Western Kenyan farm households: System dynamics model analysis. *Environ. Sci. Tech.* **45**, 3687-3694 (2011).

571 [48] Bouwman, A. F., Boumans, L. J. M. & Batjes, N. H. Emissions of N₂O and NO from
572 fertilized fields: summary of available measurement data. *Global Biogeochem. Cy.* **16**, 1058
573 (2002).

574 [49] Robertson, G. P. & Vitousek, P. M. Nitrogen in agriculture: balancing the cost of an
575 essential resource. *Annu. Rev. Env. Resour.* **34**, 97-125 (2009).

576 [50] Akiyama, H., Yan, X. Y. & Yagi, K. Evaluation of effectiveness of enhanced-efficiency
577 fertilizers as mitigation options for N₂O and NO emissions from agricultural soils: meta-
578 analysis. *Glob. Change Biol.* **16**, 1837-1846 (2010).

579 [51] Aguilera, E., Lassaletta, L., Sanz Cobeña, A., Garnier, J. & Vallejo Garcia, A. The potential
580 of organic fertilizers and water management to reduce N₂O emissions in Mediterranean
581 climate cropping systems. A review. *Agr. Ecosyst. Environ.* **164**, 32-52 (2013).

582 [52] van Kessel, C., *et al.* Climate, duration, and N placement determine N₂O emissions in
583 reduced tillage systems: a meta-analysis. *Glob. Change Biol.* **19**, 33-44 (2013).

584 [53] Millar, N., Robertson, G. P., Grace, P. R., Gehl, R. J. & Hoben, J. P. Nitrogen fertilizer
585 management for nitrous oxide (N₂O) mitigation in intensive corn (Maize) production: an
586 emissions reduction protocol for US Midwest agriculture. *Mitig. Adapt. Strat. Gl.* **15**, 185-
587 204 (2010).

588 [54] Robertson, G. P. Nitrogen-climate interactions in US agriculture. *Biogeochemistry* **114**, 41-
589 70 (2013).

590 [55] Reay, D. S. *et al.* Global agriculture and nitrous oxide emissions. *Nature Clim. Change* **2**,
591 410-416 (2012).

- 592 [56] Conrad, R. The global methane cycle: recent advances in understanding the microbial
593 processes involved. *Environ. Microbiol. Reports* **1**, 285-292 (2009).
- 594 [57] Le Mer, J. & Roger, P. Production, oxidation, emission and consumption of methane by
595 soils: A review. *Eur. J Soil Biol.* **37**, 25-50 (2001).
- 596 [58] Segers, R. Methane production and methane consumption: a review of processes underlying
597 wetland methane fluxes. *Biogeochemistry* **41**, 23-51 (1998).
- 598 [59] Suwanwaree, P. & Robertson, G.P. Methane oxidation in forest, successional, and no-till
599 agricultural ecosystems: effects of nitrogen and soil disturbance. *Soil Sci. Soc. Am. J.* **69**,
600 1722-1729 (2005).
- 601 [60] Linquist, B. A., Anders, M. M., Adviento-Borbe, M. A. A., Chaney, R. L., Nalley, L. L., da
602 Rosa, E. F.F. and van Kessel, C. Reducing greenhouse gas emissions, water use, and grain
603 arsenic levels in rice systems. *Glob Change Biol.* **21**, 407–417 (2015).
- 604 [61] Liu, Y. *et al.* Carbon Dioxide Flux from Rice Paddy Soils in Central China: Effects of
605 Intermittent Flooding and Draining Cycles. *PLOS ONE* **8**(2), e56562 (2013).
- 606 [62] FAOSTAT, <http://faostat3.fao.org/> Accessed May, 2015.
- 607 [63] van Groenigen, K. J., van Kessel, C. & Hungate, B. A. Increased greenhouse-gas intensity
608 of rice production under future atmospheric conditions. *Nature Clim. Change* **3**, 288-291
609 (2013).
- 610 [64] Levine, U. *et al.* Agriculture's impact on microbial diversity and associated fluxes of carbon
611 dioxide and methane. *The ISME Journal* **5**, 1683-1691 (2011).

- 612 [65] Su, J. *et al.* Expression of barley SUSIBA2 transcription factor yields high-starch low-
613 methane rice. *Nature* **523**, 602-606 (2015).
- 614 [66] Woolf, D., Amonette, J. E., Street-Perrott, F. A., Lehmann, J. & Joseph, S. Sustainable
615 biochar to mitigate global climate change. *Nature Communications* **1**,
616 doi:10.1038/ncomms1053 (2010). **This paper calculates the global technical potential of**
617 **greenhouse gas emission reductions by biochar systems as a function of its effects on soil**
618 **improvement.**
- 619 [67] Lynch, J. P. & Wojciechowski, T. Opportunities and challenges in the subsoil: pathways to
620 deeper rooted crops. *Journal of Experimental Botany* **66**, 2199-2210 (2015). **This paper**
621 **looks at the potential for breeding plants with more root production and tolerance to**
622 **‘problem soils’, to increase yields, nutrient and water capture and C sequestration.**
- 623 [68] Smith, S. & De Smet, I. Root system architecture: insights from Arabidopsis and cereal
624 crops Introduction. *Phil. Trans.R. Soc. B: Biol. Sci* **367**, 1441-1452 (2012).
- 625 [69] Conant, R. T., Ogle, S. M., Paul, E. A. & Paustian, K. Measuring and monitoring soil
626 organic carbon stocks in agricultural lands for climate mitigation. *Front Ecol. Environ.* **9**,
627 169-173 (2011).
- 628 [70] Paustian, K., Ogle, S. M. & Conant, R. T. Quantification and Decision Support Tools for
629 US Agricultural Soil Carbon Sequestration, in *Handbook of Climate Change and*
630 *Agroecosystems: Impact, Adaptation and Mitigation* (eds Hillel, D. & Rosenzweig, C.) 307-
631 341 (Imperial College Press, London, 2011.)

- 632 [71] Hillier, J. G. A farm-focused calculator for emissions from crop and livestock production.
 633 *Environ. Modell. Softw.* **26**, 1070-1078 (2011). On-line tool at: <http://www.coolfarmtool.org/>
 634 (Accessed May 2015).
- 635 [72] Swallow, B. M. & Goddard, T. W. Value chains for bio-carbon sequestration services:
 636 Lessons from contrasting cases in Canada, Kenya and Mozambique. *Land Use Policy* **31**, 81-
 637 89 (2013).
- 638 [73] National Academy of Sciences. *Verifying Greenhouse Gas Emissions: Methods to Support*
 639 *International Climate Agreements*. (Committee: Pacala, S., et al.). (National Academies
 640 Press, Washington, D.C.) 110 pp. (2010).
- 641 [74] Paustian, K. *et al.* COMET 2.0 – Decision support system for agricultural greenhouse gas
 642 accounting, in *Managing Agricultural Greenhouse Gases: Coordinated Agricultural*
 643 *Research through GraceNet to Address Our Changing Climate* (eds Liebig, M.,
 644 Franzluebbers, A., & Follett, R.) 251-270 (Academic Press, San Diego, CA, 2012). On-line
 645 tool at: <http://cometfarm.nrel.colostate.edu> (Accessed May 2015).
- 646 [75] Paustian, K. Bridging the data gap: engaging developing country farmers in greenhouse gas
 647 accounting. *Environ. Res. Lett.* **8**(2) (2013). **This paper proposes the development of**
 648 **mobile apps to ‘crowd-source’ local-scale knowledge of land use and management data**
 649 **to help improve GHG inventories and project-scale accounting in developing countries.**
- 650 [76] Ogle, S. M., *et al.* Scale and uncertainty in modeled soil organic carbon stock changes for
 651 US croplands using a process-based model. *Glob. Change Biol.* **16**, 810-820 (2010).

652 [77] Smith, P., *et al.* Towards an integrated global framework to assess the impacts of land use
653 and management change on soil carbon: current capability and future vision. *Glob. Change*
654 *Biol.* **18**, 2089-2101 (2012).

655 [78] Bellassen, V., *et al.* Monitoring, reporting and verifying emissions in the climate economy.
656 *Nature Clim. Change* **5**(4), 319-328 (2015).

657 [79] van Wesemael, B., *et al.* How can soil monitoring networks be used to improve predictions
658 of organic carbon pool dynamics and CO₂ fluxes in agricultural soils? *Plant Soil* **338**, 247-
659 259 (2011).

660 [80] Ogle, S. M., Breidt, F. J., Easter, M., Williams, S. & Paustian, K. An empirically based
661 approach for estimating uncertainty associated with modeling carbon sequestration in soils.
662 *Ecol. Model.* **205**, 453-463 (2007).

663 [81] Peters, W., *et al.* An atmospheric perspective on North American carbon dioxide exchange:
664 CarbonTracker. *Proc. Natl. Acad. Sci. USA* **104**, 18925–30 (2007).

665 [82] Schuh, A. E., *et al.* Evaluating atmospheric CO₂ inversions at multiple scales over highly-
666 inventoried agricultural landscape. *Glob. Change Biol.* **19**, 1424-1439 (2013).

667 [83] Miles, N. L., *et al.* Large amplitude spatial and temporal gradients in atmospheric boundary
668 layer CO₂ mole fractions detected with a tower-based network in the U.S. upper Midwest. *J.*
669 *Geophys. Res.* **117**, G01019 (2012).

670 [84] Lauvaux, T., *et al.* Constraining the CO₂ budget of the corn belt: exploring uncertainties
671 from the assumptions in a mesoscale inverse system. *Atmos. Chem. Phys.* **12**, 337-354
672 (2012).

- 673 [85] Cooley, D., Breidt, F. J., Ogle, S. M., Schuh, A. E. & Lauvaux, T. A constrained least-
674 squares approach to combine bottom-up and top-down CO₂ flux estimates. *Environ. Ecol.*
675 *Stat.* **20**, 129-146. (2013). **One of the first studies using independent ground- and**
676 **atmospheric-based methods to validate GHG emissions estimates for a large**
677 **agriculturally-intensive region.**
- 678 [86] Kadygrov, N., *et al.* Role of simulated GOSAT total column CO₂ observations in surface
679 CO₂ flux uncertainty reduction. *J. Geophys. Res.* **114**, D21208 (2009).
- 680 [87] Basu, S. *et al.* Global CO₂ fluxes estimated from GOSAT retrievals of total column CO₂.
681 *Atmos. Chem. Phys.* **13**, 8695-8717 (2013).
- 682 [88] Pacala, S. & Socolow, R. Stabilization wedges: Solving the climate problem for the next 50
683 years with current technologies. *Science* **305**, 968-972 (2004).
- 684 [89] Alexander, P., Paustian, K., Smith, P. & Moran, D. The economics of soil C sequestration
685 and agricultural emissions abatement. *Soil* **1**, 331-339 (2015).
- 686 [90] Examples of large-scale research networks conducting soil GHG flux and soil
687 measurements in geographically distributed field experiment, using uniform protocols,
688 advanced instrumentation and data portals are NitroEurope (<http://www.nitroeurope.eu>) in
689 Europe and Gracenet
690 (http://www.ars.usda.gov/research/programs/programs.htm?np_code=212&docid=21223) in
691 the US (Accessed May 2015).
- 692 [91] Dhawale, N. M. *et al.* Proximal soil sensing of soil texture and organic matter with a
693 prototype portable mid-infrared spectrometer. *Eur. J. Soil Sci.* **66**, 661-669 (2015).

694 [92] Yeluripati, J., et al. Global Research Alliance Modelling Platform (GRAMP): An open web
695 platform for modelling greenhouse gas emissions from agro-ecosystems. *Comp. & Elec.*
696 *Agric.* **111**, 112–120 (2015).

697 [93] Sanchez, P. Digital soil map of the world. *Science* **325**, 680-681 (2009).

698 [94] The Web Soil Survey (<http://websoilsurvey.sc.egov.usda.gov/App/WebSoilSurvey.aspx>) is
699 an on-line web service that provides free access to fine spatial-scale (ca. 1:15,000 – 1:20,000)
700 soil maps for the US. (Accessed May 2015).

701 [95] Awana, M. I., van Oort, P. A. J., Ahmad, R., Bastiaans, L. & Meinke, H. Farmers' views on
702 the future prospects of aerobic rice culture in Pakistan. *Land Use Policy* **42**, 517-526 (2015).

703 [96] Smith, P. *et al.* Policy and technological constraints to implementation of greenhouse gas
704 mitigation options in agriculture. *Agr. Ecosyst. Environ.* **118**, 6-28 (2007).

705 [97] Louwagie, G., S.H., Sammeth, F. & Ratinger, T. The potential of European Union policies
706 to address soil degradation in agriculture. *Land Degrad. Develop.* **22**, 5-17 (2011).

707 [98] http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=OJ:JOL_2013_165_R_0080_01
708 (Accessed May 2015).

709 [99] Examples of initial efforts, involving agricultural producers and product distributors and
710 retailers, include the Field to Market® alliance (<https://www.fieldtomarket.org/>) and the
711 Unilever Sustainable Agriculture Code ([http://www.unilever.com/Images/Unilever-](http://www.unilever.com/Images/Unilever-Sustainable-Agriculture-Code-2015_tcm244-422949.pdf)
712 [Sustainable-Agriculture-Code-2015_tcm244-422949.pdf](http://www.unilever.com/Images/Unilever-Sustainable-Agriculture-Code-2015_tcm244-422949.pdf)) (Accessed May 2015).

713 [100] Horowitz, J. K. & Just, R. E. Economics of additionality for environmental services from
714 agriculture. *J. Environ. Econ. Manag.* **66**, 105-122 (2013).

Figure text.

Fig. 1. A potential decision-tree ordering management practices towards creating GHG mitigating cropland (rice not included). For degraded, marginal lands (top of diagram) the most productive mitigation option is conversion to perennial vegetation either left unmanaged or sustainably harvested to offset fossil energy use (cellulosic biofuels). For more arable lands, multiple options could be implemented sequentially or in combination, depending on management objectives, cost and other constraints. Practices shown (see text for more discussion) are roughly arrayed from lower cost/higher feasibility options towards more costly interventions (bottom of figure). However, low cost options in one region may be a higher cost/less feasible option in another region. All options require a region-specific full-cost carbon accounting (GHG life cycle analysis) that includes potential indirect land use effects in order to define specific mitigation potentials. *Relative costs, provided as examples based on a developed region such as North America and a less developed region such as sub-Saharan Africa. †Denotes potential for major co-benefits as non-GHG ecosystem services. ‡Potential constraints that might limit or preclude practice adoption as well as potential increases in other GHGs as a consequence of practice adoption.

Fig. 2. Global potential for agricultural-based GHG mitigation, relating average per ha net GHG reduction rates and potential area (in Mha) of adoption (note log-scales). Unless otherwise noted, estimates are from Smith et al.¹⁸ based on cropland and grassland area projections for 2030. Ranges in total Pg CO₂eq yr⁻¹ represent varying adoption rates as a function of C pricing (\$20, \$50, and \$100 per Mg CO₂eq), to a maximum technical potential, i.e., full implementation of practices on the available land base. Multiple practices are aggregated for

cropland (e.g. improved crop rotations and nutrient management, reduced tillage) and grazing land (e.g., grazing management, nutrient and fire management, species introduction) categories. Practices that increase net soil C stocks and/or reduce emissions of N₂O and CH₄ are combined in each practice category. The portion of projected mitigation from C stock increase (ca. 90% of the total technical potential) would have a limited time span of 20-30 years, whereas non-CO₂ emission reduction could, in principle, continue indefinitely¹⁸. Estimates for biochar application from Woolf et al.⁶⁶ represent a technical potential only, but based on a full life cycle analysis applicable over a 100 year time span. Although global estimates of the potential impact of enhanced root phenotypes for crops have not been published, a first-order estimate of ~1 Pg CO₂eq yr⁻¹ is shown, using as an analog, global average C accrual rates (0.23 Mg C ha⁻¹ yr⁻¹) for cover crops²⁴, applied to 50% of the cropland land area used by Smith et al.¹⁸.

Fig. 3. Expanding the role for agricultural soil GHG mitigation will require an integrated research support and implementation platform. Targeted basic research on soil processes (a few examples of priority areas shown here), expanding measurement/monitoring networks and further developing global geospatial soils data can improve predictive models and reduce uncertainties. Ongoing advances in information technology and complex system and ‘Big Data’ integration, offer the potential to engage a broad-range of stakeholders, including land managers, to ‘crowd-source’ local knowledge of agricultural management practices through web-based computer and mobile apps, and help drive advanced model-based GHG metrics. This will facilitate implementation of climate-smart soil management policies, via cap-and-trade systems, product supply chain initiatives for ‘low-carbon’ consumer products, national and international

760 GHG mitigation policies and also promote more sustainable and climate-resilient agricultural
761 systems, globally.

762

[BOX 1]

Implementation strategies for soil GHG mitigation

Incentivizing farmers to adopt alternative practices that mitigate GHGs can take a variety of forms, including,

1) Regulation/taxation: Direct regulatory measures to reduce soil GHGs at the entity scale are likely politically unfeasible and costly. Taxation of N fertilizer, already used in parts of the US and Europe to reduce nitrate pollution, could function as an indirect tax to reduce N₂O emissions.

2) Subsidies: Targeted government payments/subsidies for implementing GHG-reducing practices is emerging as a policy alternative. For example, US Dept. of Agriculture programs are including GHG mitigation as a conservation goal and provisions in the EU Common Agricultural Policy link subsidy payments to ‘cross compliance’ measures that include maintenance of soil organic matter stocks⁹⁷. A more direct link to soil GHG emissions follows from a recent decision to include cropland and grassland in EU commitments under the Kyoto Protocol⁹⁸.

3) Supply chain initiatives: Major food distributors are targeting sustainability metrics, including low GHG footprints, as a consumer marketing strategy⁹⁹, setting performance standards for contracted agricultural producers, including requiring field-scale monitoring of production practices and quantification of GHG emissions.

4) Cap and trade (C&T): In a C&T system, emitters are subject to an overall emissions level or ‘cap’, in which permitted emissions decrease over time. Emitters can stay below the capped levels by reducing their own emissions and/or by purchasing surplus permits from capped entities that have exceeded their required reductions. Both compliance and voluntary markets can function as C&T systems¹⁰⁰. Within many C&T systems, a limited amount of emission

786 reductions (termed ‘offsets’) can be provided by non-capped entities. Inclusion of agricultural
787 activities as offset providers has been growing, particularly within voluntary markets. To
788 maintain the integrity of emission caps, key criteria for offset providers include demonstrating
789 *additionality*, i.e., insuring that reductions result from project interventions and not simply
790 business-as-usual trends, avoiding *leakage*, i.e., unintended emission increases elsewhere as a
791 consequence of the project activities, and providing for *permanence* (e.g., that increased soil C
792 storage, credited as a CO₂ removal, is maintained long-term).

793 **[End BOX 1]**

794

Cropland characteristics		Mitigation practices	Practice Co-benefits	Relative expense [†]		Constraints/caveats [‡]
				Developed	Less Developed	
Degraded or marginal land?	yes →	Convert to perennial set-aside or cellulosic biofuel	↓ soil erosion ↑ biodiversity ↑ water quality	\$\$	\$\$	Alternate land/livelihood for subsistence farmers Opportunity cost of removing land base Potential for leakage (i.e., land use change impact)
↓ no						
Drained, cropped organic (histosol) soils?	yes →	Restore to wetland	↑ biodiversity ↑ water quality	\$\$\$	\$\$\$	High opportunity costs of lost crop production Potential ↑ CH ₄ emissions Potential for leakage (i.e., land use change impact)
↓ no						
Severe nutrient deficiency?	yes →	Nutrient additions, liming, N-fixing species	↑ food security ↑ water quality	\$	\$\$	Availability/access to fertilizer Potential ↑ N ₂ O emissions
↓ no						
Extensive bare fallow?	yes →	Cover crops, reduced fallow vegetated fallow	↓ soil erosion ↑ water quality ↑ soil health ↑ food security	\$	\$\$	Limited applicability in dry areas
↓ no						
Excess N fertilizer use?	yes →	Reduce to economic-optimal rates	↑ water quality	\$	\$	Risk of crop production loss
↓ no						
Intensive tillage?	yes →	Reduced till, no-tillage, residue retention	↓ soil erosion ↑ soil health	\$	\$\$	Limited applicability in cold climates Potential increased equipment cost Increased herbicide use
↓ no						
Suboptimal N management?	yes →	Improve timing, placement; enhanced efficiency fertilizer	↑ water quality	\$\$	\$\$\$	Availability/access to enhanced efficiency fertilizer
↓ no						
Low residue crops?	yes →	Perennials in rotation, agroforestry, high C input species, cover crops	↑ biodiversity ↑ soil health ↑ food security	\$\$\$	\$\$	Less applicability in dry areas, shallow soils Potential opportunity costs of lost crop production
↓ no						
Available exogenous organic amendments?	yes →	Add amendments (e.g. compost, biochar)	↑ soil health ↑ food security	\$\$\$	\$\$	Dependent on lifecycle emissions of producing the amendment
↓ no						
High capacity GHG mitigation on cropland						



